

# IIT-JEE-Mathematics-Screening-2005

## SCREENING

---

1. The area of the equilateral triangle, in which three coins of radius 1 cm are placed, as shown in the figure, is:

- (a)  $6 + 4\sqrt{3}$
- (b)  $4\sqrt{3} - 6$
- (c)  $7 + 4\sqrt{3}$
- (d)  $4\sqrt{3}$

2. The area bounded by the curves  $y = (x - 1)^2$ ,  $y = (x + 1)^2$  and  $y = 1/4$  is:

- (a)  $1/3$
- (b)  $2/3$
- (c)  $1/4$
- (d)  $1/5$

3. The value of  $\int_{-2}^0 [x^3 + 3x^2 + 3x + 3 + (x+1)\cos(x+1)] dx$  is:

- (a) 0
- (b) 3
- (c) 4
- (d) 1

4. The tangent at (1, 7) to the curve  $x^2 = y - 6$  touches the circle  $x^2 + y^2 + 16x + 12y + c = 0$  at :

- (a) (6, 7)
- (b) (-6, 7)
- (c) (6, -7)
- (d) (-6, -7)

5. If  $dy/dx = xy/(x^2 + y^2)$ ,  $y(1) = 1$ , then one of the values of  $x_0$  satisfying  $y(x_0) = e$  is given by

- (a)  $e\sqrt{2}$
- (b)  $e\sqrt{3}$
- (c)  $e\sqrt{5}$
- (d)  $e/\sqrt{2}$

6. The locus of the centre of circle which touches  $(y - 1)^2 + x^2 = 1$  externally also touches x axis is:

- (a)  $x^2 = 4y$  È (0, y),  $y < 0$
- (b)  $x^2 = y$

- (c)  $y = 4x^2$   
 (d)  $y^2 = 4x \in (0, y), y \in \mathbb{R}$

7. If  $\int_{\sin x}^1 t^2 f(t) dt = 1 - \sin x \forall x \in [0, \pi/2]$  then  $f(1/\sqrt{3})$  is:

- (a) 3  
 (b)  $\sqrt{3}$   
 (c)  $1/3$   
 (d) none of these

8.

$$\binom{30}{0} \binom{30}{10} - \binom{30}{1} \binom{30}{11} + \dots + \binom{30}{20} \binom{30}{30} =$$

- (a)  ${}^{30}C_{11}$   
 (b)  ${}^{60}C_{10}$   
 (c)  ${}^{30}C_{10}$   
 (d)  ${}^{65}C_{55}$

9. A variable plane  $x/a + y/b + z/c = 1$  at a unit distance from origin cuts the coordinate axes at A, B and C. Centroid  $(x, y, z)$  satisfies the equation  $1/x^2 + 1/y^2 + 1/z^2 = K$ . The value of K is :

- (a) 9  
 (b) 3  
 (c)  $1/9$   
 (d)  $1/3$

10. Let  $f(x) = ax^2 + bx + c$ ,  $a \neq 0$  and  $D = b^2 - 4ac$ . If  $a + b$ ,  $a^2 + b^2$  and  $a^3 + b^3$  are in G.P., then :

- (a)  $D \neq 0$   
 (b)  $bD \neq 0$   
 (c)  $cD \neq 0$   
 (d)  $bc \neq 0$

11. Tangent at a point of the ellipse  $x^2/a^2 + y^2/b^2 = 1$  is drawn which cuts the coordinate axes at A and B. The minimum area of the triangle OAB is (O being the origin) :

- (a)  $ab$   
 (b)  $(a^3 + ab + b^3)/3$   
 (c)  $a^2 + b^2$   
 (d)  $((a^2 + b^2))/4$

12. A fair die is rolled. The probability that the first time 1 occurs at the even throw is :

- (a)  $1/6$

- (b) 5/11
- (c) 6/11
- (d) 5/36

**13.** If  $x dy = y (dx + y dy)$ ,  $y(1) = 1$  and  $y(x) > 0$ . Then  $y(-3) = :$

- (a) 3
- (b) 2
- (c) 1
- (d) 0

14.  $f(x) = \begin{cases} x, & \text{if } x \text{ is rational} \\ 0, & \text{if } x \text{ is irrational} \end{cases}$  and

$g(x) = \begin{cases} 0, & \text{if } x \text{ is rational} \\ x, & \text{if } x \text{ is irrational.} \end{cases}$  Then  $f - g$  is:

- (a) one-one and into
- (b) neither one-one nor onto
- (c) many one and onto
- (d) one-one and onto

**15.** A rectangle with sides  $(2n - 1)$  and  $(2m - 1)$  is divided into squares of unit length. The number of rectangle which can be formed with sides of odd length is :

- (a)  $m^2 n^2$
- (b)  $mn(m + 1)(n + 1)$
- (c)  $4^{m+n-1}$
- (d) none of these

**16.** The minimum value of  $|a + bw + cw^2|$ , where  $a, b$  and  $c$  are all not equal integers and  $w(= \omega)$  is a cube root of unity, is:

- (a)  $\sqrt{3}$
- (b)  $1/3$
- (c) 1
- (d) 0

17. If  $P = \begin{bmatrix} \sqrt{3}/2 & 1/2 \\ -1/2 & \sqrt{3}/2 \end{bmatrix}$ ,  $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$  and  $Q = PAP^T$ , then  $P^T Q^{2005} P$  is:

- (a)  $\begin{bmatrix} 1 & 2005 \\ 0 & 1 \end{bmatrix}$   
 (b)  $\begin{bmatrix} 1 & 2005 \\ 2005 & 1 \end{bmatrix}$   
 (c)  $\begin{bmatrix} 1 & 0 \\ 2005 & 1 \end{bmatrix}$   
 (d)  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

18. The shaded region, where  $P \equiv (-1, 0)$ ,  $Q \equiv (-1 + \sqrt{2}, \sqrt{2})$ ,  $R \equiv (-1 + \sqrt{2}, -\sqrt{2})$ ,  $S \equiv (1, 0)$  is represented by:

- (a)  $|z + 1| > 2$ ,  $|\arg(z + 1)| < \pi/4$   
 (b)  $|z + 1| < 2$ ,  $|\arg(z + 1)| < \pi/2$   
 (c)  $|z - 1| > 2$ ,  $|\arg(z + 1)| > \pi/4$   
 (d)  $|z - 1| < 2$ ,  $|\arg(z + 1)| > \pi/2$

19. The number of ordered pairs  $(\alpha, \beta)$ , where  $\alpha, \beta \in (-\pi, \pi)$  satisfying  $\cos(\alpha - \beta) = 1$  and  $\cos(\alpha + \beta) = 1/e$  is:

- (a) 0  
 (b) 1  
 (c) 2  
 (d) 4

20. Let  $f(x) = |x| - 1$ , then points where  $f(x)$  is not differentiable is/(are):

- (a) 0, + 1  
 (b) + 1  
 (c) 0  
 (d) 1

21. The second degree polynomial  $f(x)$ , satisfying  $f(0) = 0$ ,  $f(1) = 1$ ,  $f'(x) > 0$  for all  $x \in (0, 1)$ :

- (a)  $f(x) = x$   
 (b)  $f(x) = ax + (1 - a)x^2$ ;  $\forall a \in (0, \infty)$   
 (c)  $f(x) = ax + (1 - a)x^2$ ;  $\forall a \in (0, 2)$   
 (d) no such polynomial

22. If  $f$  is a differentiable function satisfying  $f(1/n) = 0$  for all  $n > 1$ ,  $n \in \mathbb{I}$ , then:

- (a)  $f(x) = 0$ ,  $x \in (0, 1]$   
 (b)  $f'(0) = 0 = f(0)$

- (c)  $f(0) = 0$  but  $f'(0)$  not necessarily zero  
 (d)  $|f(x)| < 1, x \in (0, 1]$

23. If  $A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & -2 & 4 \end{bmatrix}$ ,

$6A^{-1} = A^2 + cA + dI$ , then  $(c, d)$  is:

- (a)  $(-6, 11)$   
 (b)  $(-11, 6)$   
 (c)  $(11, 6)$   
 (d)  $(6, 11)$

24. In a  $\Delta ABC$ , among the following which one is true?

- (a)  $(b + c) \cos A/2 = a \sin ((B+C)/2)$   
 (b)  $(b + c) \cos ((B+C)/2) = a \sin A/2$   
 (c)  $(b - c) \cos ((B-C)/2) = a \cos (A/2)$   
 (d)  $(b - c) \cos A/2 = a \cos ((B-C)/2)$

25. If  $\vec{a}, \vec{b}, \vec{c}$  are three non zero, non coplanar vectors and  $\vec{b}_1 = \vec{b} - \frac{\vec{b} \cdot \vec{a}}{|\vec{a}|^2} \vec{a}$ ,  
 $\vec{b}_2 = \vec{b} + \frac{\vec{b} \cdot \vec{a}}{|\vec{a}|^2} \vec{a}$ . And  $\vec{c}_1 = \vec{c} - \frac{\vec{c} \cdot \vec{a}}{|\vec{a}|^2} \vec{a} - \frac{\vec{c} \cdot \vec{b}}{|\vec{b}|^2} \vec{b}$ ,  $\vec{c}_2 = \vec{c} - \frac{\vec{c} \cdot \vec{a}}{|\vec{a}|^2} \vec{a} - \frac{\vec{c} \cdot \vec{b}}{|\vec{b}_1|^2} \vec{b}_1$ ,  
 $\vec{c}_3 = \vec{c} - \frac{\vec{c} \cdot \vec{a}}{|\vec{a}|^2} \vec{a} - \frac{\vec{c} \cdot \vec{b}}{|\vec{b}_2|^2} \vec{b}_2$ ,  $\vec{c}_4 = \vec{a} - \frac{\vec{c} \cdot \vec{a}}{|\vec{a}|^2} \vec{a}$ .

Then which of the following is a set of mutually orthogonal vectors:

- (a)  $(\vec{a}, \vec{b}_1, \vec{c}_1)$   
 (b)  $(\vec{a}, \vec{b}_1, \vec{c}_2)$   
 (c)  $(\vec{a}, \vec{b}_2, \vec{c}_3)$   
 (d)  $(\vec{a}, \vec{b}_2, \vec{c}_4)$

26. If  $y = f(x)$  and  $y \cos x + x \cos y = \pi$ , then the value of  $f'(0)$  is :

- (a)  $\pi$   
 (b)  $-\pi$   
 (c)  $0$   
 (d)  $2\pi$

27. Let  $f$  be twice differentiable function satisfying  $f(1) = 1, f(2) = 4, f(3) = 9$ , then :

- (a)  $f'(x) = 2, \forall x \in (1, 3)$   
 (b)  $f'(x) = 5 = f''(x)$ , for some  $x \in (1, 3)$   
 (c) There exists at least one  $x \in (1, 3)$  such that  $f'(x) = 2$

(d) none of these

**28.** If  $X$  and  $Y$  are two non-empty sets where  $f : X \rightarrow Y$  is a function defined such that

$f(C) = \{f(x) : x \in C\}$  for  $C \subseteq X$

and  $f^{-1}(D) = \{x : f(x) \in D\}$  for  $D \subseteq Y$ ,

for any  $A \subseteq X$  and  $B \subseteq Y$  then :

(a)  $f^{-1}(f(A)) = A$

(b)  $f^{-1}(f(A)) = A$  only if  $f(X) = Y$

(c)  $f(f^{-1}(B)) = B$  only if  $B \subseteq f(X)$

(d)  $f(f^{-1}(B)) = B$