AIPMT Preliminary

CBSE PM/PD 2009

PHYSICS

1. In the nuclear decay given below:

 ${}^{A}_{Z}X \longrightarrow {}^{A}_{Z+1}Y \longrightarrow {}^{A-4}_{Z-1}B^* \longrightarrow {}^{A-4}_{Z-1}B$, the particles emitted in the sequence are:

(1) γ , β , α

(2) β, γ, α

(3) α , β , γ

(4) β, α, γ

Sol: ${}^{A}_{Z}X \longrightarrow {}^{A}_{Z+1}Y: \beta, \; {}^{A}_{Z+1}Y \longrightarrow {}^{A-4}_{Z-1}B^{*}: \alpha, \; {}^{A-4}_{Z-1}B^{*} \longrightarrow {}^{A-4}_{Z-1}B: \gamma$ (β, α, γ)

:. Correct choice: (4)

- 2. A thin circular ring of mass M and radius R is rotating in a horizontal plane about an axis vertical to its plane with a constant angular velocity ω . If two objects each of mass m be attached gently to the opposite ends of a diameter of the ring, the ring will then rotate with an angular velocity:
 - $(1) \ \frac{\omega M}{M + 2m}$

(2) $\frac{\omega(M + 2m)}{M}$

(3) $\frac{\omega M}{M + m}$

 $(4) \frac{\omega(M-2m)}{M+2m}$

Sol: $I_1 \omega_1 = I_2 \omega_2$, $I_1 = MR^2$, $I_2 = MR^2 + 2 mR^2$

$$\therefore \ \omega_2 = \frac{\mathrm{I}_1}{\mathrm{I}_2} \ \omega = \frac{\mathrm{M}}{\mathrm{M} + 2\mathrm{m}} \ \omega.$$

∴ Correct choice : (1)

- 3. In thermodynamic processes which of the following statements is not true?
 - (1) In an isochoric process pressure remains constant
 - (2) In an isothermal process the temperature remains constant
 - (3) In an adiabatic process $PV^{\gamma} = constant$
 - (4) In an adiabatic process the system is insulated from the surroundings

Sol: Pressure constant: - isobaric, not isochoric

- 4. The number of photo electrons emitted for light of a frequency v (higher than the threshold frequency $\boldsymbol{\nu}_0)$ is proportional to:
 - (1) Threshold frequency (v_0)
- (2) Intensity of light
- (3) Frequency of light (v)
- **(4)** $v v_0$

Sol: Saturation current ∞ intensity

∴ Correct choice : (2)

- 5. A simple pendulum performs simple harmonic motion about x = 0 with an amplitude a and time period T. The speed of the pendulum at $x = \frac{a}{2}$ will be:
 - (1) $\frac{\pi a}{T}$

 $(2) \frac{3\pi^2 a}{T}$

(3) $\frac{\pi a \sqrt{3}}{T}$


 $(4) \frac{\pi a \sqrt{3}}{2T}$

Sol: Speed $v = \omega \sqrt{a^2 - x^2}$, $x = \frac{a}{2}$


$$\therefore \mathbf{v} = \omega \sqrt{\mathbf{a}^2 - \frac{\mathbf{a}^2}{4}} = \omega \sqrt{\frac{3\mathbf{a}^2}{4}}$$
$$= \frac{2\pi}{T} \frac{\mathbf{a}\sqrt{3}}{2} = \frac{\pi \mathbf{a}\sqrt{3}}{T}$$

:. Correct choice: (3)

6. See the electric circuit shown in this Figure. Which of the following equations is a correct equation for it?



- (1) $\varepsilon_2 i_2 r_2 \varepsilon_1 i_1 r_1 = 0$
- (2) $\varepsilon_2 (i_1 + i_2) R + i_2 r_2 = 0$
- (3) $\varepsilon_1 (i_1 + i_2) R + i_1 r_1 = 0$ (4) $\varepsilon_1 (i_1 + i_2) R i_1 r_1 = 0$



 $\varepsilon_1 - (i_1 + i_2)R - i_1 r_1 = 0.$

∴ Correct choice : (4)

- 7. A body, under the action of a force $\vec{F}=6\ \hat{i}\ -8\ \hat{j}\ +10\ \hat{k}$, acquires an acceleration of 1 m/s². The mass of this body must be:
 - (1) 10 kg
- (2) 20 kg
- (3) $10\sqrt{2} \text{ kg}$
 - (4) $2\sqrt{10} \text{ kg}$

Sol: $\vec{F} = 6\hat{i} - 8\hat{j} + 10\hat{k}$,

$$|F| = \sqrt{36 + 64 + 100} = 10\sqrt{2} N$$

$$a = 1 \, \mathrm{ms}^{-2}$$

$$\therefore m = \frac{10\sqrt{2}}{1} = 10\sqrt{2} \text{ kg}$$

∴ Correct choice : (3)

8. The symbolic representation of four logic gates are given below:



The logic symbols for OR, NOT and NAND gates are respectively:

- (1) (iv), (i), (iii)
- (2) (iv), (ii), (i)
- (3) (i), (iii), (iv)
- **(4)** (iii), (iv), (ii)

9. If \vec{F} is the force acting on a particle having position vector \vec{r} and $\vec{\tau}$ be the torque of this force about the origin, then:

(1)
$$\vec{r} \cdot \vec{\tau} > 0$$
 and $\vec{F} \cdot \vec{\tau} < 0$

(2)
$$\vec{r} \cdot \vec{\tau} = 0$$
 and $\vec{F} \cdot \vec{\tau} = 0$

(3)
$$\vec{r} \cdot \vec{\tau} = 0$$
 and $\vec{F} \cdot \vec{\tau} \neq 0$

(4)
$$\vec{r} \cdot \vec{\tau} \neq 0$$
 and $\vec{F} \cdot \vec{\tau} = 0$

Sol:
$$\vec{\tau} = \vec{r} \times \vec{F} \Rightarrow \vec{r} \cdot \vec{\tau} = 0$$
 $\vec{F} \cdot \vec{\tau} = 0$

∴ Correct choice : (2)

10. The two ends of a rod of length L and a uniform cross-sectional area A are kept at two temperatures T_1 and T_2 ($T_1 > T_2$). The rate of heat transfer, $\frac{dQ}{dt}$ through the rod in a steady state is given by:

(1)
$$\frac{dQ}{dt} = \frac{k(T_1 - T_2)}{LA}$$

(2)
$$\frac{dQ}{dt} = k L A (T_1 - T_2)$$

(3)
$$\frac{dQ}{dt} = \frac{k A (T_1 - T_2)}{L}$$

(4)
$$\frac{dQ}{dt} = \frac{kL(T_1 - T_2)}{A}$$

$$\mbox{Sol:} \ \, \frac{dQ}{dt} = \frac{kA(T_1 \, - \, T_2)}{L} \label{eq:Sol}$$

∴ Correct choice : (3)

- 11. A p-n photodiode is fabricated from a semiconductor with a band gap of 2.5 eV. It can detect a signal of wavelength:
 - (1) 4000 nm
- (2) 6000 nm
- (3) 4000 Å
- (4) 6000 Å

Sol:
$$\lambda_{max} = \frac{hc}{E} = \frac{6.6 \times 10^{-34} \times 3 \times 10^{8}}{2.5 \times 1.6 \times 10^{-19}} \simeq 5000 \text{ Å}$$

$$\lambda < \lambda_{\text{max}} = 4000 \text{ Å}$$

: Correct choice: (3)

- 12. If the dimensions of a physical quantity are given by $M^a \ L^b \ T^c$, then the physical quantity will be:
 - (1) Velocity if a = 1, b = 0, c = -1
 - (2) Acceleration if a = 1, b = 1, c = -2
 - (3) Force if a = 0, b = -1, c = -2
 - (4) Pressure if a = 1, b = -1, c = -2

Sol: Pressure = $\frac{\text{MLT}^{-2}}{\text{L}^2}$ = $\text{ML}^{-1}\text{T}^{-2}$

$$\Rightarrow$$
 a = 1, b = -1, c = -2.

∴ Correct choice : (4)

- 13. A transistor is operated in common-emitter configuration at V_c = 2 V such that a change in the base current from 100 μA to 200 μA produces a change in the collector current from 5 mA to 10 mA. The current gain is:
 - (1) 100
- **(2)** 150
- **(3)** 50
- (4) 75

Sol: $\Delta I_E = \Delta I_B + \Delta I_C$

$$\beta = \frac{\Delta I_{C}}{\Delta I_{B}}$$

$$\Delta I_{\rm C} = 5 \times 10^{-3} \,\mathrm{A}$$

$$\Delta I_{\rm B} = 100 \times 10^{-6} \, \rm A$$

$$\beta = \frac{5}{100} \times 1000 = 50$$

∴ Correct choice : (3)

- 14. The mass of a lift is 2000 kg. When the tension in the supporting cable is 28000 N, then its acceleration is:
 - (1) 4 ms^{-2} upwards.

(2) 4 ms^{-2} downwards.

(3) $14 \text{ ms}^{-2} \text{ upwards.}$

(4) 30 ms^{-2} downwards.

Sol: 2000 a = 28000 - 20000 = 8000

$$a = \frac{8000}{2000} = 4 \text{ ms}^{-2} \uparrow$$

- 15. Four identical thin rods each of mass M and length ℓ , form a square frame. Moment of inertia of this frame about an axis through the centre of the square and perpendicular to its plane is:

- (1) $\frac{2}{3} \text{ M} \ell^2$ (2) $\frac{13}{3} \text{ M} \ell^2$ (3) $\frac{1}{3} \text{ M} \ell^2$ (4) $\frac{4}{3} \text{ M} \ell^2$

Sol: $\frac{\text{mL}^2}{12} + \frac{\text{mL}^2}{4} = \frac{4\text{mL}^2}{12} = \frac{\text{mL}^2}{3}$

Total M.I. = $4 \times \frac{\text{mL}^2}{2}$

: Correct choice: (4)

16. Each of the two strings of length 51.6 cm and 49.1 cm are tensioned separately by 20 N force. Mass per unit length of both the strings is same and equal to 1 g/m. When both the strings vibrate simultaneously the number of beats is:

(1) 7

- **(2)** 8
- (3) 3

(4) 5

Sol: $f_1 = \frac{1}{2\ell_1} \sqrt{\frac{T}{m}}, f_2 = \frac{1}{2\ell_2} \sqrt{\frac{T}{m}}, f_2 - f_1 = \frac{1}{2} \sqrt{\frac{T}{m} \frac{(\ell_1 - \ell_2)}{\ell_1 \ell_2}}$

$$\sqrt{\frac{T}{m}} = \sqrt{\frac{20}{10^{-3}}} = \sqrt{2} \times 10^2 = 1.414 \times 100 = 141.4$$

$$\frac{\ell_1 - \ell_2}{\ell_1 \ell_2} = \frac{(51.6 - 49.1) \times 10^2}{51.6 \times 49.1} = \frac{2.5 \times 10^2}{50 \times 50} = \frac{1}{10}$$

$$\therefore f_2 - f_1 = \frac{1}{2} \times 141.4 \times \frac{1}{10} = 7 \text{ beats}$$

:. Correct choice: (1)

- 17. The number of beta particles emitted by a radioactive substance is twice the number of alpha particles emitted by it. The resulting daughter is an:
 - (1) isomer of parent


(2) isotone of parent

(3) isotope of parent

(4) isobar of parent

:. Correct choice: (3)

18. The Figure shows a plot of photo current versus anode potential for a photo sensitive surface for three different radiations. Which one of the following is a correct statement?




Retarding potential

Anode potential

- (1) curves (a) and (b) represent incident radiations of same frequency but of different intensities.
- (2) curves (b) and (c) represent incident radiations of different frequencies and different intensities.
- (3) curves (b) and (c) represent incident radiations of same frequency having same intensity.
- (4) curves (a) and (b) represent incident radiations of different frequencies and different intensities.

∴ Correct choice : (1)

19. The Figure shows elliptical orbit of a planet m about the sum S. The shaded area SCD is twice the shaded area SAB. If t₁ is the time for the planet of move from C to D and t₂ is the time to move from A to B then:



(1)
$$t_1 = 4t_2$$

(1)
$$t_1 = 4t_2$$
 (2) $t_1 = 2t_2$

(3)
$$t_1 = t_2$$

(4)
$$t_1 > t_2$$

Sol: $SCD : A_1 - t_1$ (areal velocity constant)

$$SAB:A_2-t_2$$

$$\frac{A_1}{t_1} = \frac{A_2}{t_2}, t_1 = t_2 \cdot \frac{A_1}{A_2}, A_1 = 2A_2$$

$$\therefore t_1 = 2t_2$$

:. Correct choice: (2)

- 20. A black body at 227°C radiates heat at the rate of 7 Cals/cm²s. At a temperature of 727°C, the rate of heat radiated in the same units will be:
 - (1) 50
- **(2)** 112
- (3) 80
- **(4)** 60

Sol: $E = \sigma T^4$,

$$\frac{E_2}{E_1} = \left(\frac{T_2}{T_1}\right)^4 = \left(\frac{1000}{500}\right)^4 = 16$$


$$T_1 = 500 \text{ K}$$

$$T_2 = 1000 \text{ K}$$

$$\therefore E_2 = 16 \times 7 = 112 \text{ cal/cm}^2 \text{ s.}$$

:. Correct choice : (2)


- 21. The driver of a car travelling with speed 30 m/sec towards a hill sounds a horn of frequency 600 Hz. If the velocity of sound in air is 330 m/s, the frequency of reflected sound as heard by driver is:
 - (1) 555.5 Hz
- (2) 720 Hz
- (3) 500 Hz
- (4) 550 Hz




$$f' = \frac{v}{v - 30} f$$
, $f'' = \frac{v + 30}{v} f' = \frac{v + 30}{v - 30} f = \frac{360}{300} \times 600$
= 720 Hz


:. Correct choice : (2)

22. A wire of resistance 12 ohms per meter is bent to form a complete circle of radius 10 cm. The resistance between its two diametrically opposite points, A and B as shown in the Figure, is:



- (1) 3Ω
- (2) $6 \pi \Omega$
- (3) 6Ω
- (4) $0.6 \pi \Omega$





$$2\pi R \longrightarrow 12 \Omega$$

$$\therefore R = \frac{6 \times 6}{12} = 3 \Omega$$

:. Correct choice: (1)

- 23. A rectangular, a square, a circular and an elliptical loop, all in the (x-y) plane, are moving out of a uniform magnetic field with a constant velocity, $\overrightarrow{V}=v\,\widehat{i}$. The magnetic field is directed along the negative z axis direction. The induced emf, during the passage of these loops, out of the field region, will not remain constant for:
 - (1) the circular and the elliptical loops.
 - (2) only the elliptical loop.
 - (3) any of the four loops.
 - (4) the rectangular, circular and elliptical loops.
- **Sol:** As the loop leaves the magnetic field, area in magnetic field decreases for all loops, so induced emf does not remain constant. (Any of four loops)

∴ Correct choice : (3)

- 24. A galvanometer having a coil resistance of 60Ω shows full scale deflection when a current of 1.0 amp passes through it. It can be converted into an ammeter to read currents upto 5.0 amp by:
 - (1) putting in series a resistance of 15 Ω
 - (2) putting in series a resistance of 240 Ω
 - (3) putting in parallel a resistance of 15 Ω
 - (4) putting in parallel a resistance of 240 Ω

Sol:
$$G = 60 \Omega$$
, $I_g = 1.0 A$, $I = 5 A$.

$$I_g G = (I - I_g) S$$
,

$$S = \frac{I_g G}{I - I_g} = \frac{1}{5 - 1} \times 60 = 15 \Omega$$

putting 15 Ω in parallel.

25. Power dissipated in an LCR series circuit connected to an a.c source of emf ϵ is:

(1)
$$\frac{\epsilon^2 \sqrt{R^2 + \left(Lw - \frac{1}{Cw}\right)^2}}{R}$$

(2)
$$\frac{\varepsilon^2 \left[R^2 + \left(Lw - \frac{1}{Cw} \right)^2 \right]}{R}$$

(3)
$$\frac{\epsilon^2 R}{\sqrt{R^2 + \left(Lw - \frac{1}{Cw}\right)^2}}$$

(4)
$$\frac{\varepsilon^2 R}{\left[R^2 + \left(Lw - \frac{1}{Cw}\right)^2\right]}$$

Sol: Power dissipated in series LCR:
$$P = I^2R = \frac{\epsilon^2}{|Z|^2} R = \frac{\epsilon^2 R}{\left[R^2 + \left(\omega L - \frac{1}{\omega C}\right)^2\right]}$$

∴ Correct choice : (4)

26. Three concentric spherical shells have radii a, b and c (a < b < c) and have surface charge densities σ , – σ and σ respectively. If V_A , V_B and V_C denote the potentials of the three shells, then for c = a + b, we have:

(1)
$$V_C = V_B \neq V_A$$

(2)
$$V_C \neq V_B \neq V_A$$


(3)
$$V_C = V_B = V_A$$

(4)
$$V_C = V_A \neq V_B$$

Sol: c = a + b.

$$\boldsymbol{v}_{A} = \frac{\sigma Q}{\epsilon_{0}} - \frac{\sigma \boldsymbol{b}}{\epsilon_{0}} + \frac{\sigma \boldsymbol{c}}{\epsilon_{0}} = \frac{\sigma}{\epsilon_{0}} \left[\boldsymbol{c} - (\boldsymbol{b} - \boldsymbol{a}) \right]$$

$$\begin{split} \boldsymbol{V}_{B} &= \frac{-\ \sigma \boldsymbol{b}}{\epsilon_{0}} + \frac{1}{4\pi\epsilon_{0}} \cdot \frac{\sigma \times 4\pi\boldsymbol{a}^{2}}{\boldsymbol{b}} + \frac{\sigma \boldsymbol{c}}{\epsilon_{0}} \\ &= \frac{\sigma}{\epsilon_{0}} \left[\boldsymbol{c} - \frac{(\boldsymbol{b}^{2} - \boldsymbol{a}^{2})}{\boldsymbol{b}} \right] \end{split}$$



$$\begin{split} V_C &= \frac{\sigma c}{\epsilon_0} - \frac{1}{4\pi\epsilon_0} \cdot \frac{\sigma \times 4\pi b^2}{c} + \frac{1}{4\pi\epsilon_0} \cdot \frac{\sigma \times 4\pi a^2}{c} = \frac{\sigma}{\epsilon_0} \Bigg[c - \frac{(b^2 - a^2)}{c} \Bigg] \\ &= \frac{\sigma}{\epsilon_0} \Big[c - (b - a) \Big] \end{split}$$

$$\mathbf{V}_{\mathrm{A}} = \mathbf{V}_{\mathrm{C}} \neq \mathbf{V}_{\mathrm{B}}$$

27.	a velocity v and		unit length of the wa	Water leaves the hose with ater jet. What is the rate at
	(1) mv ²	(2) $\frac{1}{2}$ mv ²	(3) $\frac{1}{2}$ m ² v ²	(4) $\frac{1}{2}$ mv ³
Sol:	m : mass per ui	nit length		
	∴ rate of mass	$per sec = \frac{mx}{t} = mv$		
	Rate of K.E. =	$\frac{1}{2} (mv) v^2 = \frac{1}{2} mv^3$	3	
				∴ Correct choice : (4)
28.	A bar magnet	having a magnetic	moment of 2×10^4	JT ⁻¹ is free to rotate in a
	The work done	e. A horizontal mag in taking the magne from the field is:	metic field $B = 6 \times 4$ slowly from a dire	10 ⁻⁴ T exists in the space. ction parallel to the field to
	(1) 12 J	(2) 6 J	(3) 2 J	(4) 0.6 J
Sol:	Work done = M	$B(\cos\theta_1 - \cos\theta_2)$		
	$= nB\left(1 - \frac{1}{2}\right)$	$=\frac{2\times10^4\times6\times}{2}$	$\frac{10^{-4}}{} = 6 \text{ J}$	
				∴ Correct choice : (2)
29.	approaches a t		arge \mathbf{z}_2 and mass 1	e of charge \mathbf{z}_1 and mass \mathbf{M}_1 \mathbf{M}_2 , the distance of closest
	(1) directly pro	portional to $\mathbf{z}_1\mathbf{z}_2$		
	(2) inversely p	roportional to z ₁		
	(3) directly pro	oportional to mass M	1	
	(4) directly pro	portional to $M_1 \times M$	2	
				∴ Correct choice : (1)
30.	The power em		number of photons	ed by a helium neon laser. s arriving per sec. On the
	(1) 3×10^{16}	(2) 9×10^{15}	(3) 3×10^{19}	(4) 9×10^{17}

Sol: $\lambda = 667 \times 10^{-9} \text{ m}, P = 9 \times 10^{-3} \text{ W}$

 $P = \frac{Nhc}{\lambda}$, N: No. of photons emitted/sec.

$$N = \frac{9 \times 10^{-3} \times 667 \times 10^{-9}}{6.6 \times 10^{-34} \times 3 \times 10^{8}}$$
$$= \frac{9 \times 6.67 \times 10^{-10}}{3 \times 6.6 \times 10^{-26}} \approx 3 \times 10^{16} / \text{sec}$$

∴ Correct choice : (1)

- 31. A wave in a string has an amplitude of 2 cm. The wave travels in the + ve direction of x axis with a speed of 128 m/sec. and it is noted that 5 complete waves fit in 4 m length of the string. The equation describing the wave is:
 - (1) y = (0.02) m sin (15.7x 2010t)
 - (2) y = (0.02) m sin (15.7x + 2010t)
 - (3) y = (0.02) m sin (7.85x 1005t)
 - (4) y = (0.02) m sin (7.85x + 1005t)

Sol:
$$A = 2 \text{ cm}, \frac{\omega}{k} = 128 \text{ ms}^{-1}, 5\lambda = 4, \lambda = \frac{4}{5} \text{ m}$$

 $y = A \sin(kx - \omega t)$,

$$k = \frac{2\pi}{\lambda} = \frac{2\pi \times 5}{4} = \frac{31.4}{4} = 7.85$$

$$y = 0.02 \text{ m sin} (7.857 - 1005 \text{ t})$$

$$\omega = 128 \times 7.85 = 1005$$

:. Correct choice: (3)

- **32.** Which one of the following equations of motion represents simple harmonic motion?
 - (1) acceleration = -k(x + a)
 - (2) acceleration = k(x + a)
 - (3) acceleration = kx
 - (4) acceleration = $-k_0x + k_1x^2$

Where k, k_0 , k_1 and a are all positive.

Sol: a = -kX, X = x + a.

	internal resistance r) as a function of the current (I) flowing through it. The slop and intercept, of the graph between V and I, then, respectively, equal:				
	(1) – r and \in	(2) r and − ∈	(3) $- \in$ and r	$(4) \in \text{and} - r$	
Sol:	V + ir = E				
	$V = V_A - V_B$				
	E-ir				
	$\frac{\partial V}{\partial i} = - r, i = 0,$	v = E			
	\therefore slope = -r, inte	rcept = E			
			÷.	Correct choice : (1)	
34.	If a diamagnetic s magnet, it is:	ubstance is brought	near the north or the	e south pole of a bar	
	(1) repelled by the	e north pole and attra	acted by the south pole)	
	(2) attracted by the	ne north pole and rep	elled by the south pole		
	(3) attracted by be	oth the poles			
	(4) repelled by both	th the poles			
			∴ ·	Correct choice : (4)	
35.	overtake the bus i		on a straight road. As at a distance of 1 km hase the bus?		
	(1) 40 ms^{-1}	(2) 25 ms ⁻¹	(3) 10 ms^{-1}	(4) 20 ms^{-1}	
Sol:	Let v be the relativ	ve velocity of scooter	w.r.t b as		
	$v = v_S - v_B$				
	$\therefore v_S = v + v_B, v =$	$=\frac{1000}{100} = 10 \text{ ms}^{-1}$	S I 1 km	$u = 10 \text{ ms}^{-1}$	
	∴ velocity of scoot	$er = 20 \text{ ms}^{-1}$			
				Correct choice : (4)	
36.	Sodium has body of The lattice parameters		tance between two ne	arest atoms is 3.7 Å.	
	(1) 4.3 Å	(2) 3.0 Å	(3) 8.6 Å	(4) 6.8 Å	

33. A student measures the terminal potential difference (V) of a cell (of emf \in and

Sol:
$$3.7 = \frac{\sqrt{3}}{2}$$
 a

$$a = \frac{2 \times 3.7}{\sqrt{3}} = 4.3 \text{ Å}$$

:. Correct choice: (1)

- 37. The internal energy change in a system that has absorbed 2 Kcals of heat and done 500 J of work is:
 - (1) 6400 J
- (2) 5400 J
- (3) 7900 J
- (4) 8900 J

Sol: $Q = \Delta U + W$

$$\Delta U = Q - W = 2 \times 4.2 \times 1000 - 500 = 8400 - 500$$

$$= 7900 J$$

:. Correct choice: (3)

- 38. Three capacitors each of capacitance C and of breakdown voltage V are joined in series. The capacitance and breakdown voltage of the combination will be:

 - (1) 3C, $\frac{V}{3}$ (2) $\frac{C}{3}$, 3V (3) 3C, 3V
- (4) $\frac{C}{3}$, $\frac{V}{3}$

Sol: Q = CV

$$V_{\text{eff}} = V + V + V = 3 V$$

$$\frac{1}{C_{\text{eff}}} = \frac{1}{C} + \frac{1}{C} + \frac{1}{C} \Rightarrow C_{\text{eff}} = \frac{C}{3}$$

$$\left(\frac{\mathrm{C}}{3}, 3\mathrm{V}\right)$$

:. Correct choice : (2)


- 39. An explosion blows a rock into three parts. Two parts go off at right angles to each other. These two are, 1 kg first part moving with a velocity of 12 ms⁻¹ and 2 kg second part moving with a velocity of 8 ms⁻¹. If the third part flies off with a velocity of $4~\mathrm{ms}^{-1}$, its mass would be:
 - (1) 7 kg
- (2) 17 kg
- (3) 3 kg
- (4) 5 kg

Sol: My cos $\theta = 12$

Mv $\sin \theta = 16$

$$\tan\theta = \frac{16}{12} = \frac{4}{3}$$

$$M = \frac{12 \times 5}{4 \times 3} = \frac{60}{12} = 5 \text{ kg}$$



:. Correct choice: (4)

40. A particle starts its motion from rest under the action of a constant force. If the distance covered in first 10 seconds is \mathbf{S}_1 and that covered in the first 20 seconds is \mathbf{S}_2 , then:

(1)
$$S_2 = 3S_1$$

(2)
$$S_2 = 4S_1$$

(3)
$$S_2 = S_1$$

(4)
$$S_2 = 2S_1$$

Sol:
$$s_1 = \frac{1}{2} a \times t_1^2$$
, $s_2 = \frac{1}{2} a \times t_2^2$

$$\therefore \frac{s_1}{s_2} = \left(\frac{t_1}{t_2}\right)^2 = \left(\frac{10}{20}\right)^2 = \frac{1}{4}$$

$$s_2 = 4 s_1$$

∴ Correct choice : (2)

41. A body of mass 1 kg is thrown upwards with a velocity 20 m/s. It momentarily comes to rest after attaining a height of 18 m. How much energy is lost due to air friction? ($g = 10 \text{ m/s}^2$)

Sol:
$$\frac{1}{2} \text{ mv}^2 - \text{mgh} = \frac{1}{2} \times 1 \times 400 - 1 \times 18 \times 10$$

$$= 200 - 180 = 20 J$$

:. Correct choice: (4)

42. A conducting circular loop is placed in a uniform magnetic field 0.04 T with its plane perpendicular to the magnetic field. The radius of the loop starts shrinking at 2 mm/s. The induced emf in the loop when the radius is 2 cm is:

(1)
$$4.8 \pi \mu V$$

(2)
$$0.8 \pi \mu V$$

(3)
$$1.6 \pi \mu V$$

(4)
$$3.2 \pi \mu V$$

Sol:
$$e = -B \frac{d}{dt} (\pi r^2) = -B \pi 2r \frac{dr}{dt}$$

$$\begin{aligned} r &= 2 \text{ cm, e} = -0.04 \times 3.14 \times 2 \times 2 \times 10^{-2} \times 2 \times 10^{-3} = -0.04 \times 25.12 \times 10^{-7} \\ &= 100.48 \times 10^{-7} \\ &= 32 \ \pi \times 10^{-7} \\ &= 3.2 \ \pi \times 10^{-6} \ \text{V} = 3.2 \ \pi \ \text{u V} \end{aligned}$$

: Correct choice: (4)

43. The magnetic force acting on a charged particle of charge – 2 μC in a magnetic field of 2T acting in y direction, when the particle velocity is

$$(2\hat{i} + 3\hat{j}) \times 10^6 \text{ ms}^{-1}$$
, is:

(1) 4 N is z direction

(2) 8 N is y direction

(3) 8 N in z direction

(4) 8 N in - z direction

Sol:
$$\vec{F} = q (\vec{V} \times \vec{B}) = -2 \times 10^{-6} \text{ C} [2 \times 2 \times 10^{6}] = -8 \text{ N z-axis}$$

: Correct choice: (4)

44. Two bodies of mass 1 kg and 3 kg have position vectors \hat{i} + $2\hat{j}$ + \hat{k} and $-3\hat{i}-2\hat{j}+\hat{k}$, respectively. The centre of mass of this system has a position

(1)
$$-2\hat{i} - \hat{j} + \hat{k}$$
 (2) $2\hat{i} - \hat{j} - 2\hat{k}$ (3) $-\hat{i} + \hat{j} + \hat{k}$ (4) $-2\hat{i} + 2\hat{k}$

$$(3) - \hat{i} + \hat{j} + \hat{k}$$

(4)
$$-2\hat{i} + 2\hat{l}$$

Sol:
$$\vec{R} = \frac{\vec{m_1} \cdot \vec{k_1} + \vec{m_2} \cdot \vec{k_2}}{(\vec{m_1} + \vec{m_2})} = \frac{1}{4} \left[-8\hat{i} - 4\hat{j} + 4\hat{k} \right] = -2\hat{i} - \hat{j} + \hat{k}$$

∴ Correct choice : (1)

45. The electric potential at a point (x, y, z) is given by $V = -x^2y - xz^3 + 4$ The electric field \overrightarrow{E} at that point is:

(1)
$$\vec{E} = \hat{i} (2xy + \hat{i} (x^2 + y^2) + \hat{k} (3xz - y^2)$$

(2)
$$\vec{E} = \hat{i}z^3 + \hat{j}xyz + \hat{k}z^2$$

(3)
$$\vec{E} = \hat{i} (2xy - z^3) + \hat{j} xu^2 + \hat{k} 3z^2 x$$

(4)
$$\vec{E} = \hat{i} (2xy + z^3) + \hat{i}x^2 + \hat{k} 3xz^2$$

Sol:
$$\vec{E} = -\frac{\partial V}{\partial r} = \left[-\frac{\partial V}{\partial x} \hat{i} - \frac{\partial V}{\partial y} \hat{j} - \frac{\partial V}{\partial z} \hat{k} \right]$$
$$= \left[(2 xy + z^3) \hat{i} + \hat{j} x^2 + \hat{k} 3 xz^2 \right]$$

: Correct choice: (4)


- **46.** The mean free path of electrons in a metal is 4×10^{-8} m. The electric field which can given on an average 2 eV energy to an electron in the metal will be in units of V/m:
 - (1) 5×10^{-11} (2) 8×10^{-11} (3) 5×10^{7} (4) 8×10^{7}

Sol:
$$E = \frac{V}{d} = \frac{2}{4 \times 10^{-8}} = 0.5 \times 10^8 = 5 \times 10^7 \text{ Vm}^{-1}$$

:. Correct choice: (3)

- 47. The ionization energy of the electron in the hydrogen atom in its ground state is 13.6 eV. The atoms are excited to higher energy levels to emit radiations of 6 wavelengths. Maximum wavelength of emitted radiation corresponds to the transition between:
 - (1) n = 3 to n = 1 states
- (2) n = 2 to n = 1 states
- (3) n = 4 to n = 3 states
- (4) n = 3 to n = 2 states

Sol:
$$\frac{n(n-1)}{2} = 6$$



$$n^2 - n - 12 = 0$$

$$(n-4)(n+3) = 0$$
 or $n=4$

: Correct choice: (3)

- **48.** Under the influence of a uniform magnetic field, a charged particle moves with constant speed V in a circle of radius R. The time period of rotation of the particle:
 - (1) depends on R and not on V
 - (2) is independent of both V and R
 - (3) depends on both V and R
 - (4) depends on V and not on R

Sol:
$$T = \frac{2\pi m}{qB}$$

: Correct choice: (2)

49. The electric field part of an electromagnetic wave in a medium is represented by $\mathbf{E}_{\mathbf{x}} = \mathbf{0};$

$$E_{y} = 2.5 \frac{N}{C} \cos \left[\left(2\pi \times 10^{6} \frac{\text{rad}}{\text{m}} \right) t - \left(\pi \times 10^{-2} \frac{\text{rad}}{\text{s}} \right) x \right];$$

 $E_z = 0$. The wave is:

- (1) moving along x direction with frequency 10^6 Hz and wave length 100 m.
- (2) moving along x direction with frequency 10^6 Hz and wave length 200 m.
- (3) moving along x direction with frequency 10^6 Hz and wave length 200 m.
- (4) moving along y direction with frequency $2\pi\times 10^6$ Hz and wave length 200 m.

Sol:
$$E_v = E_0 \cos(\omega t - kx)$$

$$\omega = 2 \pi f = 2 \pi \times 10^6$$
 $\therefore f = 10^6 \text{ Hz}$

$$\frac{2\pi}{\lambda} = k = \pi \times 10^{-2} \text{ m}^{-1}, \ \lambda = 200 \text{ m}$$

∴ Correct choice : (2)

- **50.** A block of mass M is attached to the lower end of a vertical spring. The spring is hung from a ceiling and has force constant value k. The mass is released from rest with the spring initially unstretched. The maximum extension produced in the length of the spring will be:
 - (1) 2 Mg/k
- (2) 4 Mg/k
- (3) Mg/2k
- (4) Mg/k

Sol: ka = mg


$$a = \frac{mg}{k}$$

:. Correct choice : (4)

BIOLOGY

- **51.** Which one of the following is **correct** pairing of a body part and the kind of muscle tissue that moves it?
 - (1) Biceps of upper arm Smooth muscle fibres
 - (2) Abdominal wall Smooth muscle
 - (3) Iris Involuntary smooth muscle
 - (4) Heart wall Involuntary unstriated muscle
- ∴ Correct choice : (2)
- **52.** The epithelial tissue present on the inner surface of bronchioles and fallopian tubes is:
 - (1) Glandular
- (2) Ciliated
- (3) Squamous
- (4) Cuboidal
- ∴ Correct choice : (2)


53. Study the pedigree chart given below:



What does it show?

- (1) Inheritance of a condition like phenylketonuria as an autosomal recessive trait
- (2) The pedigree chart is wrong as this is not possible
- (3) Inheritance of a recessive sex-linked disease like haemophilia
- (4) Inheritance of a sex-linked inborn error of metabolism like phenylketonuria

Sol:



Parents needs to be heterozygous as two of their children are known to be sufferer of the disease. It cannot be recessive sex-linked inheritance because then the male parent would also be sufferer.

54.	Manganese is required in:		
	(1) Plant cell wall formation		
	(2) Photolysis of water during photosyr	nthesis	
	(3) Chlorophyll synthesis		
	(4) Nucleic acid synthesis		
		.:	Correct choice : (2)
55 .	Polyethylene glycol method is used for:		
	(1) Biodiesel production	(2) Seedless fruit pr	roduction
	(3) Energy production from sewage	(4) Gene transfer w	rithout a vector
		:	Correct choice : (4)
56.	The floral formula \bigoplus \circlearrowleft $K_{(5)} \stackrel{\longleftarrow}{C_{(5)}} A_5 \stackrel{\frown}{C_{(5)}} A_5 \stackrel{\frown}{C_{(5)$	$\frac{4}{5}(2)$ is that of:	
	(1) Soybean (2) Sunnhemp	(3) Tobacco	(4) Tulip
Sol:	Soyabean and Sunnhemp have mono flower and perianth.	carpellary pistil and	d tulip has trimerous
			Correct choice : (3)
57 .	Which one of the following groups o triploblastic?	f animals is bilater	ally symmetrical and
	(1) Aschelminthes (round worms)	(2) Ctenophores	
	(3) Sponges	(4) Coelenterates (C	Cnidarians)
		:	Correct choice : (1)
58.	Which one of the following is commonly plants?	y used in transfer of	foreign DNA into crop
	(1) Meloidogyne incognita	(2) Agrobacterium	n tumefaciens
	(3) Penicillium expansum	(4) Trichoderma l	narzianum
		:	Correct choice : (2)
59.	Which one of the following is the corremenstrual cycle?	ct matching of the e	vents occurring during
	(1) Proliferative phase	: Rapid regeneration	n of myometrium and
		maturation of Gra	afian follicle.
	(2) Development of corpus luteum	: Secretory phase a of progesterone.	nd increased secretion


: breakdown of myometrium and ovum (3) Menstruation not fertilised. (4) Ovulation : LH and FSH attain peak level and sharp fall in the secretion of progesterone. ∴ Correct choice : (2) **60.** Which one is the wrong pairing for the disease and its causal organism? (1) Black rust of wheat - Puccinia graminis (2) Loose smut of wheat - Ustilago nuda (3) Root-knot of vegetables - Meloidogyne sp (4) Late blight of potato - Alternaria solani ∴ Correct choice : (4) 61. Global agreement in specific control strategies to reduce the release of ozone depleting substances, was adopted by: (1) The Montreal Protocol (2) The Koyoto Protocol (3) The Vienna Convention (4) Rio de Janeiro Conference ∴ Correct choice : (1) **62**. What is **true** about Bt toxin? (1) Bt protein exists as active toxin in the Bacillus. (2) The activated toxin enters the ovaries of the pest to sterilise it and thus prevent its multiplication. (3) The concerned Bacillus has antitoxins. (4) The inactive protoxin gets converted into active form in the insect gut. ∴ Correct choice : (4) **63**. **Peripatus** is a connecting link between: (1) Mollusca and Echinodermata (2) Annelida and Arthropoda (3) Coelenterata and Porifera (4) Ctenophora and Platyhelminthis ∴ Correct choice : (2) 64. T.O. Diener discovered a: (1) Free infectious DNA (2) Infectious protein (4) Free infectious RNA (3) Bacteriophage Sol: T.O. Diener discovered viroid which is free infectious RNA. ∴ Correct choice : (4)

65.	5. Seminal plasma in humans is rich in:				
	(1) fructose and calcium but has no enzymes				
	(2) glucose and certain enzymes but has no calcium				
	(3) fructose and o	certain enzymes but p	oor in calcium		
	(4) fructose, calci	ium and certain enzyr	nes		
				∴ Correct choice : (3)	
66.	A fruit developed	from hypanthodium	inflorescence is cal	led:	
	(1) Sorosis	(2) Syconus	(3) Caryopsis	(4) Hesperidium	
				∴ Correct choice : (2)	
67.	The cell junctions	s called tight, adherin	g and gap junction	s are found in:	
	(1) Connective tis	ssue	(2) Epithelial tiss	sue	
	(3) Neural tissue		(4) Muscular tiss	sue	
				∴ Correct choice : (2)	
68.	What will happened ?	n if the stretch recep	tors of the urinary	bladder wall are totally	
	(1) Micturition w	rill continue			
	(2) Urine will cor	ntinue to collect norm	ally in the bladder		
	(3) There will be	no micturition			
	(4) Urine will not	t collect in the bladder	ŗ		
Sol:	: Micturition is same as urination. Urination is the act of passing urine which is a reflex phenomenon. As urine accumulates in bladder the stretch receptors are activated that pass the stimulus to the spinal cord. In the absence of stretch receptors the urine would get collected and probably overflow.				
				∴ Correct choice : (3)	
69.		vorm is pricked with the fluid that comes		s outer surface without	
	(1) coelomic fluid	(2) haemolymph	(3) slimy mucus	(4) excretory fluid	
				∴ Correct choice : (1)	

70.	O. The most popularly known blood grouping is the ABO grouping. It is named ABO and not ABC, because "O" in it refers to having:				
	(1) overdominance of this type on the genes for A and B types				
	(2) one antibody only – either anti-A or	anti-B on the RB	$\mathbb{C}\mathbf{s}$		
	(3) no antigens A and B on RBCs				
	(4) other antigens besides A and B on RBCs				
			∴ Correct choice : (3)		
71.	One of the synthetic auxin is:				
	(1) IAA (2) GA	(3) IBA	(4) NAA		
			∴ Correct choice : (4)		
72.	A person likely to develop tetanus is im	munised by admir	istering:		
	(1) Preformed antibodies	(2) Wide spectrus	m antibiotics		
	(3) Weakened germs	(4) Dead germs			
Sol:	: Tetanus toxoid is a vaccine consisting of growth products of Clostridium tetani treated with formaladehyde serving as an active immunising agent. Hence is is weakened germs.				
			∴ Correct choice : (3)		
73.	Alzheimer disease in humans is associa	ted with the defici	ency of:		
	(1) glutamic acid				
	(2) acetylcholine				
	(3) gamma aminobutyric acid (GABA)				
	(4) dopamine				
			∴ Correct choice : (2)		
74.	Biochemical Oxygen Demand (BOD) in	a river water:			
	(1) has no relationship with concentrat	ion of oxygen in th	e water.		
	(2) gives a measure of salmonella in t	he water.			
	(3) increases when sewage gets mixed	with river water.			
	(4) remains unchanged when algal block	om occurs.			
			∴ Correct choice : (3)		

75.	The genetic defect – adenosine dea permanently by:	aminase (ADA) deficiency may be cured
	(1) administering adenosine deaminase	e activators.
	(2) introducing bone marrow cells prostages.	oducing ADA into cells at early embryonic
	(3) enzyme replacement therapy.	
	(4) periodic infusion of genetically e ADA cDNA.	ngineered lymphocytes having functional
		∴ Correct choice : (2)
76 .	Compared to blood our lymph has:	
	(1) plasma without proteins	(2) more WBCs and no RBCs
	(3) more RBCs and less WBCs	(4) no plasma
		∴ Correct choice : (2)
77.	Sickle cell anemia is:	
	(1) caused by substitution of valine by haemoglobin	y glutamic acid in the beta globin chain of
	(2) caused by a change in a single base	pair of DNA
	(3) characterized by elongated sickle lil	ke RBCs with a nucleus
	(4) an autosomal linked dominant trait	t
		∴ Correct choice : (2)
78.	Which of the following plant species bioethanol?	s you would select for the production of
	(1) Zea mays (2) Pongamia	(3) Jatropha (4) Brassica
		∴ Correct choice : (3)
79.	When breast feeding is replaced by less the infants below the age of one year an	s nutritive food low in proteins and calories; re likely to suffer from:
	(1) Rickets (2) Kwashiorkor	(3) Pellagra (4) Marasmus
		∴ Correct choice : (2)

80.	A young infant may be feeding entirely on mother's milk which is white in colour but the stools which the infant passes out is quite yellowish. What is this yellow colour due to?				
	(1) Bile pigment	ts passed through b	ile juice		
	(2) Undigested	milk protein casein			
	(3) Pancreatic ju	uice poured into du	odenum		
	(4) Intestinal ju	ice			
				∴ Correct choice : (1)	
81.	Which one of the	e following has max	ximum genetic dive	rsity in India?	
	(1) Mango	(2) Wheat	(3) Tea	(4) Teak	
				∴ Correct choice : (2)	
82.	Oxygenic photos	synthesis occurs in:			
	(1) Oscillatoria	a	(2) Rhodosp	irillum	
	(3) Chlorobium	n	(4) Chromat	ium	
				∴ Correct choice : (1)	
83.	There is no DNA	A in:			
	(1) Mature RBC	L's	(2) A mature	spermatozoan	
	(3) Hair root		(4) An enucle	ated ovum	
Sol:	An enucleated o	vum has DNA in m	itochondria.		
				∴ Correct choice : (1)	
84.	Given below is a	a schematic break-u	p of the phases / st	ages of cell cycle:	
			A B C Mitosis		



(1) C-Karyokinesis

(2) D-Synthetic phase

(3) A-Cytokinesis

cycle?

(4) B-Metaphase

85. Tiger is not a resident in which one of the following national park?

(1) Sunderbans

(2) Gir

(3) Jim Corbett

(4) Ranthambhor

∴ Correct choice : (2)

86. Which one of the following statements is **true** regarding digestion and absorption of food in humans?

(1) Fructose and amino acids are absorbed through intestinal mucosa with the help of carrier ions like Na⁺.

(2) Chylomicrons are small lipoprotein particles that are transported from intestine into blood capillaries.

(3) About 60% of starch is hydrolysed by salivary amylase in our mouth.

(4) Oxyntic cells in our stomach secrete the proenzyme pepsinogen.

∴ Correct choice : (1)

87. Synapsis occurs between:

(1) mRNA and ribosomes


(2) spindle fibres and centromere

(3) two homologous chromosomes

(4) a male and a female gamete

∴ Correct choice : (3)

88. Given below is a diagrammatic sketch of a portion of human male reproductive system. Select the correct set of the names of the parts labelled A, B, C, D.





A	В	\mathbf{C}	D
(1) vas deferens	seminal vesicle	prostate	bulbourethral gland
(2) vas deferens	seminal vesicle	bulbourethral gland	prostate
(3) ureter	seminal vesicle	prostate	bulbourethral gland
(4) ureter	prostate	seminal vesicle	bulbourethral gland
			:. Correct choice : (1)

89.	9. What is not true for genetic code?								
	(1) It is nearly universal								
	(2) It is degenerate								
	(3) It is unambigu	ous							
	(4) A codon in mR	NA is read i	n a non-c	ontig	uous fa	shio	n		
							∴ Correct c	hoice :	(4)
90.	Which one of the fe	ollowing pla	nts is mo	noecio	ous?				
	(1) Pinus	(2) Cycas		(3) F	Papaya		(4) Mar	chanti	a
							∴ Correct c	hoice :	(1)
91.	Cyclic photophospl	horylation r	esults in t	the fo	rmatio	n of			
	(1) ATP and NAD	PH		(2) A	ATP, N	ADP	H and ${\rm O}_2$		
	(3) ATP			(4) N	NADPE	I			
							∴ Correct c	hoice :	(3)
92.	The letter T in T-l	ymphocyte 1	refers to:						
	(1) Thalamus	(2) Tonsil		(3) T	hymu	S	(4) Thyr	oid	
					-		∴ Correct c	hoice	(3)
93.	Foetal ejection refl	lex in huma	n female i	is ind	uced by	y:			
	(1) release of oxyte	ocin from pi	tuitary						
	(2) fully developed	l foetus and	placenta						
	(3) differentiation	of mammar	y glands						
	(4) pressure exert	ed by amnio	tic fluid						
							∴ Correct c	hoice :	(2)
94.	Anatomically fair dicotyledonous ste		icotyledon	nous	root	is	distinguished	from	the
	(1) Absence of seco	ondary phloe	em	(2) I	Presenc	ce of	cortex		
	(3) Position of prot	toxylem		(4) A	Absence	e of s	secondary xylen	ı	
							:. Correct c	hoice	(3)
95 .	Plasmodesmata ar	e:							
	(1) Locomotary str	ructures							
	(2) Membranes co	nnecting the	e nucleus	with	plasma	alem	ma		
	(3) Connections be	etween adja	cent cells						
	(4) Lignified ceme	nted layers	between o	cells					
							∴ Correct c	hoice	(3)

96.	6. Removal of introns and joining the exons in a defined order in a transcription unit is called:			
	(1) Tailing	(2) Transformation	(3) Capping	(4) Splicing
				∴ Correct choice : (4)
97.	Phylogenetic syste	m of classification is	based on :	
	(1) Morphological	features	(2) Chemical con	stituents
	(3) Floral characte	ers	(4) Evolutionary	relationships
				∴ Correct choice : (4)
98.	Which part of hum	nan brain is concerne	d with the regulat	ion of body temperature?
	(1) Cerebellum		(2) Cerebrum	
	(3) Hypothalamus		(4) Medulla Oblo	ngata
				∴ Correct choice : (3)
99.	Semiconservative	replication of DNA w	as first demonstra	ted in:
	(1) Escherichia o	coli	(2) Streptococc	us pneumoniae
	(3) Salmonella ty	phimurium	(4) Drosophila	melanogaster
				∴ Correct choice : (1)
100.	Which one of the f	ollowing pairs of anim	nals comprises 'jav	wless fishes'?
	(1) Mackerals and	Rohu	(2) Lampreys and	d hag fishes
	(3) Guppies and ha	ag fishes	(4) Lampreys and	d eels
				∴ Correct choice : (2)
101.	Which of the follow	ving is a pair of viral	diseases?	
	(1) Common Cold,	AIDS	(2) Dysentery, Co	ommon Cold
	(3) Typhoid, Tuber	rculosis	(4) Ringworm, A	IDS
				∴ Correct choice : (1)
102.	Aerobic respirator	y pathway is approp	riately termed:	
	(1) Parabolic	(2) Amphibolic	(3) Anabolic	(4) Catabolic
				∴ Correct choice : (2)

103. A country with a high rate of population growth took measures to reduce it. The Figure below shows age-sex pyramids of populations A and B twenty years apart. Select the correct interpretation about them:





Interpretations:

- (1) "B" is earlier pyramid and shows stabilised growth rate.
- (2) "B" is more recent showing that population is very young.
- (3) "A" is the earlier pyramid and no change has occurred in the growth rate.
- (4) "A" is more recent and shows slight reduction in the growth rate.

∴ Correct choice : (4)

- 104. Cytoskeleton is made up of:
 - (1) Callose deposits

- (2) Cellulosic microfibrils
- (3) Proteinaceous filaments
- (4) Calcium carbonate granules
- **Sol:** Cytoskeleton is made up of microfilaments and microtubules whose major constituents are actin and tubulin respectively.

∴ Correct choice : (3)

- 105. An example of axile placentation is:
 - (1) Dianthus
- (2) Lemon
- (3) Marigold
- (4) Argemone
- ∴ Correct choice : (2)
- 106. Which one of the following has haplontic life cycle?
 - (1) Polytrichum

(2) Ustilago

(3) Wheat

(4) Funaria

107.	'. Steps taken by the Government of India to control air pollution include:			
	(1) compulsory PUC (Pollution Under Control) certification of petrol driven vehicles which tests for carbon monoxide and hydrocarbons.			
	(2) permission to u fuel for vehicles		with a maximum of	500 ppm sulphur as
	(3) use of non-polluting Compressed Natural Gas (CNG) only as fuel by all buses and trucks.			
	(4) compulsory mix diesel.	ring of 20% ethyl a	lcohol with petrol and	d 20% biodiesel with
			∴ (Correct choice : (1)
108.	Which one of the f habit?	following is conside	red important in the	development of seed
	(1) Heterospory		(2) Haplontic life cyc	le
	(3) Free-living game	etophyte	(4) Dependent sporo	phyte
			∴ (Correct choice : (1)
109.	The annular and sprotoxylem when the		nducting elements gen	erally develop in the
	(1) elongating	(2) widening	(3) differentiating	(4) maturing
			∴ (Correct choice : (4)
110.	The correct sequer	nce of plants in a hy	drosere is:	
	(1) Volvox \longrightarrow Hy	y drilla \longrightarrow Pistia	\longrightarrow Scirpus \longrightarrow La	ntana —→ Oak
	(2) Pistia \longrightarrow Vol	vox	\longrightarrow Hydrilla \longrightarrow Oal	$k \longrightarrow Lantana$
	(3) Oak → Lanta	ana→ Volvox	\rightarrow Hydrilla \longrightarrow Pist	ia→ Scirpus
	(4) Oak → Lanta	ana —→ Scirpus —	→ Pistia —→ Hydri	$lla \longrightarrow Volvox$
			∴ (Correct choice : (1)
111.	Stroma in the chlor	oplasts of higher pla	ant contains:	
	(1) Light-dependen	t reaction enzymes		
	(2) Ribosomes			
	(3) Chlorophyll			
	(4) Light-independe	ent reaction enzyme	es	
			∴ 0	Correct choice : (4)

112.	2. A health disorder that results from the deficiency of thyroxine in adults and characterised by (i) a low metabolic rate, (ii) increase in body weight and (iii) tendency to retain water in tissues is:			
	(1) simple goitre (2) myxoedema	(3) cretinism	(4) hypothyroidism	
Sol:	Deficiency of thyroxine in adults is sp by low metabolic rate, increase in boot tissues.			
			∴ Correct choice : (2)	
113.	Mannitol is the stored food in:			
	(1) Porphyra (2) Fucus	(3) Gracillaria	(4) Chara	
			∴ Correct choice : (2)	
114.	Which one of the following pairs is wro	ongly matched?		
	(1) Alcohol – nitrogenase	(2) Fruit juice –	pectinase	
	(3) Textile – amylase	(4) Detergents –	lipase	
			∴ Correct choice : (1)	
115.	Which of the following is not used as a	biopesticide?		
	(1) Trichoderma harzianum	(2) Nuclear Poly	hedrosis Virus (NPV)	
	(3) Xanthomonas campestris	(4) Bacillus thu	ıringiensis	
			∴ Correct choice : (3)	
116.	Which one of the following is a vascula	r cryptogam?		
	(1) Ginkgo (2) Marchantia	(3) Cedrus	(4) Equisetum	
			∴ Correct choice : (4)	
117.	In a standard ECG which one of representation of the respective activity			
	(1) S – start of systole	(2) T – end of dia	astole	
	(3) P – depolarisation of the atria	(4) R – repolaris	ation of ventricles	
			∴ Correct choice : (3)	
118.	Uric acid is the chief nitrogenous comp	onent of the excre	tory products of:	
	(1) Earthworm (2) Cockroach	(3) Frog	(4) Man	
			∴ Correct choice : (2)	
119.	Guard cells help in:			
	(1) Transpiration	(2) Guttation		
	(3) Fighting against infection	(4) Protection ag	gainst grazing	
			∴ Correct choice : (1)	

120.	Montreal Protocol aims at:					
	(1) Biodiversity conservation					
	(2) Control of water pollution					
	(3) Control of CO_2 emission					
	(4) Reduction of o	(4) Reduction of ozone depleting substances				
				∴ Correct choice : (4)		
121.	DDT residues ar because DDT is:	DDT residues are rapidly passed through food chain causing biomagnification because DDT is:				
	(1) moderately toxic		(2) non-toxic to aquatic animals			
	(3) water soluble		(4) lipo soluble			
				∴ Correct choice : (4)		
122.	Vegetative propag	gation in mint occurs	by:			
	(1) Offset	(2) Rhizome	(3) Sucker	(4) Runner		
				∴ Correct choice : (3)		
123.	Select the incorr	ect statement from th	ne following:			
	(1) Galactosemia is an inborn error of metabolism					
	(2) Small population size results in random genetic drift in a population					
	(3) Baldness is a sex-limited trait					
	(4) Linkage is an exception to the principle of independent assortment in heredity			nt assortment in heredity		
				∴ Correct choice : (3)		
124.	Cotyledons and te	esta respectively are e	edible parts in:			
	(1) walnut and ta	marind	(2) french bean	and coconut		
	(3) cashew nut ar	nd litchi	(4) groundnut a	nd pomegranate		
				∴ Correct choice : (4)		
125.	Which one of the	Which one of the following statements is correct?				
	(1) Benign tumours show the property of metastasis.					
	(2) Heroin accelerates body functions.					
	(3) Malignant tumours may exhibit metastasis.					
	(4) Patients who have undergone surgery are given cannabinoids to relieve pain.					
				∴ Correct choice : (3)		

126.	The correct sequesperms in a mature		netic stages lead	ing to the formation of			
	(1) spermatogonia – spermatocyte – spermatid – sperms						
	(2) spermatid – spermatocyte – spermatogonia – sperms						
	(3) spermatogonia – spermatid – spermatocyte – sperms						
	(4) spermatocyte – spermatogonia – spermatid – sperms						
				∴ Correct choice : (1)			
127.	Use of anti-histam	Use of anti-histamines and steroids give a quick relief from:					
	(1) Nausea	(2) Cough	(3) Headache	(4) Allergy			
				∴ Correct choice : (4)			
128.	Chipko movement	Chipko movement was launched for the protection of:					
	(1) Forests	(2) Livestock	(3) Wet lands	(4) Grasslands			
				∴ Correct choice : (1)			
129.	Which one of the following is the most likely root cause why menstruation is not taking place in regularly cycling human female?						
	(1) maintenance of the hypertrophical endometrial lining						
	(2) maintenance of high concentration of sex hormones in the blood stream						
	(3) retention of well-developed corpus luteum						
	(4) fertilisation of the ovum						
				∴ Correct choice : (4)			
130.	Globulins containe	bulins contained in human blood plasma are primarily involved in:					
	(1) osmotic balance of body fluids		(2) oxygen transport in the blood				
	(3) clotting of blood		(4) defence mechanisms of body				
				∴ Correct choice : (4)			
131.	Palisade parenchyma is absent in leaves of:						
	(1) Mustard	(2) Soybean	(3) Gram	(4) Sorghum			
				∴ Correct choice : (4)			
132.	In barley stem vas	cular bundles are:					
	(1) closed and scattered		(2) open and in a ring				
	(3) closed and radial		(4) open and scattered				
				∴ Correct choice : (1)			

133. Which one of the following is the **correct** matching of three items and their grouping category?

Items Group

(1) ilium, ischium, pubis – coxal bones of pelvic girdle

(2) actin, myosin, rhodopsin – muscle proteins

(3) cytosine, uracil, thiamine – pyrimidines

(4) malleus, incus, cochlea – ear ossicles

∴ Correct choice : (1)

134. Somaclones are obtained by

(1) Plant breeding (2) Irradiation

(3) Genetic engineering (4) Tissue culture

∴ Correct choice : (4)

- 135. In the case of peppered moth (Biston betularia) the black-coloured form became dominant over the light-coloured form in England during industrial revolution. This is an example of:
 - (1) appearance of the darker coloured individuals due to very poor sunlight
 - (2) protective mimicry
 - (3) inheritance of darker colour character acquired due to the darker environment
 - (4) natural selection whereby the darker forms were selected
- Sol: This is a phenomenon of industrial melanism. The moths rested during day time when their predators (birds) are active. During industrial revolution, the surrounding areas were covered with soot and hence dark forms got camouflaged. This offered protection to dark forms when coal was used. Later when electricity was source of energy the environment became lighter (absence of soot) and more of the paler forms of moth were sighted.

∴ Correct choice : (2)

- **136.** Transgenic plants are the ones:
 - (1) generated by introducing foreign DNA into a cell and regenerating a plant from that cell.
 - (2) produced after protoplast fusion in artificial medium.
 - (3) grown in artificial medium after hybridization in the field.
 - (4) produced by a somatic embryo in artificial medium.

137.	Which one of the following pairs of stomach totally undigested?	food components	in humans reaches the
	(1) Starch and fat	(2) Fat and cellul	ose
	(3) Starch and cellulose	(4) Protein and s	tarch
			∴ Correct choice : (2)
138.	A change in the amount of yolk and its	distribution in the	egg will affect:
	(1) Pattern of cleavage		
	(2) Number of blastomeres produced		
	(3) Fertilization		
	(4) Formation of zygote		
			∴ Correct choice : (1)
139.	Middle lamella is composed mainly of:		
	(1) Muramic acid	(2) Calcium pecta	ate
	(3) Phosphoglycerides	(4) Hemicellulose	?
			∴ Correct choice : (2)
140.	Elbow joint is an example of:		
	(1) hinge joint	(2) gliding joint	
	(3) ball and socket joint	(4) pivot joint	
			∴ Correct choice : (1)
141.	Which of the following is a symbiotic ni	trogen fixer ?	
	(1) Azotobacter (2) Frankia	(3) Azolla	(4) Glomus
			∴ Correct choice : (2)
142.	Whose experiments cracked the DNA and discovered unequivocally that a genetic code is a "triplet"?		
	(1) Hershey and Chase	(2) Morgan and S	Sturtevant
	(3) Beadle and Tatum	(4) Nirenberg and	d Mathaei
			∴ Correct choice : (4)
143.	Which one of the following types of org in a pond ecosystem?	anisms occupy mo	re than one trophic level
	(1) Fish (2) Zooplankton	(3) Frog	(4) Phytoplankton
Sol:	Fish could be primary consumer as well	l as secondary cons	sumer.
			∴ Correct choice : (1)

144.	Which one of the following acids is a derivative of carotenoids?					
	(1) Indole-3-acetic	acid	(2) Gibberellic acid			
	(3) Abscisic acid		(4) Indole butyrio	eacid		
				∴ Correct choice : (3)		
145.	5. The bacterium Bacillus thuringiensis is widely used in contemporary biologas:					
	(1) Insecticide					
	(2) Agent for production of dairy products					
	(3) Source of industrial enzyme					
	(4) Indicator of water pollution					
				∴ Correct choice : (1)		
146.	An example of a se	eed with endosperm, j	perisperm, and car	runcle is:		
	(1) coffee	(2) lily	(3) castor	(4) cotton		
				∴ Correct choice : (3)		
147.	Reduction in vascu	ılar tissue, mechanica	al tissue and cuticl	e is characteristic of :		
	(1) Mesophytes	(2) Epiphytes	(3) Hydrophytes	(4) Xerophytes		
				∴ Correct choice : (3)		
148.	Point mutation involves:					
	(1) Change in sing	le base pair	(2) Duplication			
	(3) Deletion		(4) Insertion			
				∴ Correct choice : (1)		
149.	Which one of the following $correctly$ describes the location of some body parts in the earthworm $Pheretima$?					
	(1) Four pairs of spermathecae in 4 – 7 segments.					
	(2) One pair of ovaries attached at intersegmental septum of $14^{\rm th}$ and $15^{\rm th}$ segments.					
	(3) Two pairs of te	stes in $10^{ ext{th}}$ and $11^{ ext{th}}$ s	egments.			
	(4) Two pairs of ac	ecessory glands in 16	– 18 segments.			
				∴ Correct choice : (3)		
150.	The kind of tissue is also found in:	that forms the suppo	rtive structure in	our pinna (external ears)		
	(1) nails	(2) ear ossicles	(3) tip of the nose	e (4) vertebrae		
				∴ Correct choice : (3)		

151. The state of hybridization of C_2 , C_3 , C_5 and C_6 of the hydrocarbon,

$$\begin{array}{c|cccc} CH_3 & CH_3 \\ | & | \\ CH_3 - C - CH = CH - CH - C \equiv CH \\ 7 & 6| & 5 & 4 & 3 & 2 & 1 \\ CH_3 & & & & & \end{array}$$

is in the following sequence:

- (1) sp^3 , sp^2 , sp^2 and sp
- (2) sp, sp 2 , sp 2 and sp 3
- (3) sp, sp 2 , sp 3 and sp 2
- (4) sp, sp 3 , sp 2 and sp 3

Sol: sp, sp^3, sp^2, sp^3

:. Correct choice: (4)

- **152.** Oxidation numbers of P in PO_4^{3-} , of S in SO_4^{2-} and that of Cr in $Cr_2O_7^{2-}$, are respectively:
 - (1) + 3, + 6 and + 5

(2) + 5, + 3 and + 6

(3) - 3, + 6 and + 6

(4) + 5, + 6 and + 6

Sol: PO_4^{3-} (P = + 5)

$$SO_4^{2-}$$
 (S = + 6)

$$Cr_2O_7^{2-}$$
 (Cr = + 6)

:. Correct choice: (4)

- **153.** Lithium metal crystallises in a body centred cubic crystal. If the length of the side of the unit cell of lithium is 351 pm, the atomic radius of the lithium will be:
 - (1) 151.8 pm
- (2) 75.5 pm
- (3) 300.5 pm
- (4) 240.8 pm

Sol: a $\sqrt{3} = 4 \text{ r}$

$$r = \frac{a\sqrt{3}}{4} = \frac{351 \times 1.732}{4} = 151.98 \text{ pm}$$

∴ Correct choice : (1)

154.	Which of the following reactions is an example of nucleophilic susbtitution reaction?					
	(1) 2 RX + 2 N	$Na \longrightarrow R - R + 2 N$	ΙaX			
	(2) RX + H ₂ -	\longrightarrow RH + HX				
	(3) RX + Mg -	\longrightarrow RMgX				
	(4) RX + KOH	$I \longrightarrow ROH + KX$				
Sol:	X is replaced	by OH				
	•			∴ Correct choice : (4)		
155.	In the case of a	ılkali metals, the co	ovalent character dec	creases in the order:		
	(1) MF > MCl >	> MBr > MI	(2) MF > MCl >	MI > MBr		
	(3) MI > MBr >	- MC1 > MF	(4) MCl > MI >	MBr > MF		
Sol:	MI > MBr > M0	Cl > MF. As the siz	e of the anion decrea	ases covalency decreases		
				∴ Correct choice : (3)		
156.	Which one of the elements with the following outer orbital configurations material exhibit the largest number of oxidation states?					
	(1) $3d^54s^1$	(2) $3d^54s^2$	(3) $3d^24s^2$	(4) $3d^34s^2$		
Sol:	The configuration $3d^5 4s^2$ can have various oxidation states upto + 7.					
				∴ Correct choice : (2)		
157.	7. The stability of + 1 oxidation state increases in the sequence:					
	(1) Tl < In < Ga	a < Al	(2) In < Tl < Ga	a < Al		
	(3) Ga < In < A	1 < T1	(4) Al < Ga < I	n < T1		
Sol:	: The order is due to 'inert pair effect'					
				∴ Correct choice : (4)		
158.	Given:					
	(i) $Cu^{2+} + 2e^{-} -$	\longrightarrow Cu, $E^0 = 0.337$	V			
	(ii) Cu ²⁺ + e ⁻ —	\longrightarrow Cu ⁺ , E ⁰ = 0.153	V			
	Electrode poter	ntial, E ^o for the rea	ection, $Cu^+ + e^- \longrightarrow$	Cu, will be:		
	(1) 0.90 V	(2) 0.30 V	(3) 0.38 V	(4) 0.52 V		

Sol:
$$Cu^{2+} + 2e^{-} \longrightarrow Cu$$
; $\Delta G^{0} = -nE^{0} F = -2 \times F \times 0.337 = -0.674 F$
 $Cu^{+} \longrightarrow Cu^{2+} + e^{-}$; $\Delta G^{0} = -nE^{0} F = -1 \times F \times -0.153 = 0.153 F$

$$Cu^{+} + e^{-} \longrightarrow Cu$$
: $\Delta G^{0} = -0.521 \text{ F} = -nE^{0}F$: $n = 1$. $E^{0} = +0.52 \text{ V}$

.: Correct choice: (4)

159. For the reaction,
$$N_2 + 3H_2 \longrightarrow 2NH_3$$
, if $\frac{d \left[NH_3 \right]}{dt} = 2 \times 10^{-4} \text{ mol L}^{-1} \text{ s}^{-1}$, the value of $\frac{-d \left[H_2 \right]}{dt}$ would be:

(1)
$$4 \times 10^{-4} \text{ mol L}^{-1} \text{ s}^{-1}$$

(2)
$$6 \times 10^{-4} \text{ mol L}^{-1} \text{ s}^{-1}$$

(3)
$$1 \times 10^{-4} \text{ mol L}^{-1} \text{ s}^{-1}$$

(4)
$$3 \times 10^{-4} \text{ mol L}^{-1} \text{ s}^{-1}$$

Sol:
$$-\frac{1}{3} \frac{d [H_2]}{dt} = \frac{1}{2} \frac{d [NH_3]}{dt}$$
$$-\frac{d [H_2]}{dt} = \frac{3}{2} \frac{d [NH_3]}{dt} = \frac{3}{2} \times 2 \times 10^{-4} = 3 \times 10^{-4} \text{ mol } L^{-1} \text{ s}^{-1}$$

:. Correct choice: (4)

160. Consider the following reaction,

$$ethanol \xrightarrow{PBr_3} X \xrightarrow{alc. \ KOH} Y \xrightarrow{(i) \ H_2SO_4 \ room \ temperature} \xrightarrow{(ii) \ H_2O, \ heat} Z;$$

the product Z is:

(1)
$$CH_3CH_2 - O - CH_2 - CH_3$$
 (2) $CH_3 - CH_2 - O - SO_3H_3$

(2)
$$CH_3 - CH_2 - O - SO_3H$$

(4)
$$CH_2 = CH_2$$

:. Correct choice: (3)

161. The energy absorbed by each molecule (A $_2$) of a substance is 4.4×10^{-19} J and bond energy per molecule is 4.0×10^{-19} J. The kinetic energy of the molecule per atom will be:

(1)
$$2.2 \times 10^{-19} \text{ J}$$

(2)
$$2.0 \times 10^{-19}$$
 J

(3)
$$4.0 \times 10^{-20} \text{ J}$$

(4)
$$2.0 \times 10^{-20}$$
 J

Sol: K.E per atom =
$$\frac{\left(4.4 \times 10^{-19}\right) - \left(4.0 \times 10^{-19}\right)}{2} = \frac{0.4 \times 10^{-19}}{2} = 2.0 \times 10^{-20}$$

∴ Correct choice : (4)

162. Amongst the elements with following electronic configurations, which one of them may have the highest ionization energy?

(1) Ne
$$[3s^23p^2]$$

(2) Ar
$$[3d^{10}4s^24p^3]$$

(3) Ne
$$[3s^23p^1]$$

(4) Ne
$$[3s^23p^3]$$

 ${f Sol:}\,\,\,{f Smallest}$ atom having half filled p-sub shell has highest ${f I_0}$ value

∴ Correct choice : (4)

163. In the reaction


 ${\rm BrO}_3^-$ (aq) + 5 ${\rm Br}_{(aq)}^-$ + 6H $^+$ o 3 ${\rm Br}_2(1)$ + 3 ${\rm H_2O}_{(1)}$. The rate of appearance of bromine (Br $_2$) is related to rate of disappearance of bromide ions as following:

(1)
$$\frac{d \left(Br_2\right)}{dt} = -\frac{5}{3} \frac{d \left(Br^-\right)}{dt}$$

(2)
$$\frac{d \left(Br_2\right)}{dt} = \frac{5}{3} \frac{d \left(Br^-\right)}{dt}$$

(3)
$$\frac{d \left(Br_2\right)}{dt} = \frac{3}{5} \frac{d \left(Br^-\right)}{dt}$$

(4)
$$\frac{d \left(Br_2\right)}{dt} = -\frac{3}{5} \frac{d \left(Br^-\right)}{dt}$$



- 169. Half life period of a first-order reaction is 1386 seconds. The specific rate constant of the reaction is:
 - (1) $0.5 \times 10^{-2} \text{ s}^{-1}$

(2) $0.5 \times 10^{-3} \text{ s}^{-1}$

(3) $5.0 \times 10^{-2} \text{ s}^{-1}$

- **(4)** $5.0 \times 10^{-3} \text{ s}^{-1}$
- **Sol:** $t_{1/2} = \frac{0.693}{k}$; $k = \frac{0.693}{1386} = 0.5 \times 10^{-3} \text{ s}^{-1}$

∴ Correct choice : (2)

- **170.** Which one of the following is employed as a tranquilizer?
 - (1) Naproxen

(2) Tetracycline

(3) Chlorpheninamine

(4) Equanil

∴ Correct choice : (4)

- 171. ${\rm Al_2O_3}$ is reduced by electrolysis at low potentials and high currents. If 4.0×10^4 amperes of current is passed through molten $\mathrm{Al_2O_3}$ for 6 hours, what mass of aluminium is produced? (Assume 100% current efficiency. At. mass of Al = 27 g mol⁻¹)
 - (1) 8.1×10^4 g

- (2) 2.4×10^5 g (3) 1.3×10^4 g (4) 9.0×10^3 g

Sol: Total current = $4.0 \times 10^4 \times 6 \times 60 \times 60$ C

96500 C liberates 9 g of Al (1 g. eq)

 $(4\times10^4\times6\times60\times60)$ C liberates 8.1×10^4 g of Al

∴ Correct choice : (1)

- **172.** Benzene reacts with CH₃Cl in the presence of anhydrous AlCl₃ to form:
 - (1) Chlorobenzene (2) Benzylchloride (3) Xylene
- (4) Toluene

$$\textbf{Sol:} \ \ \, \overbrace{\bigcirc} \ \, + \ \, \text{CH}_3\text{Cl} \ \, \underbrace{\frac{\text{Anhyd.}}{\text{AlCl}_3}} \ \, \underbrace{\bigcirc} \ \, + \ \, \text{HCl}$$

∴ Correct choice : (4)

173. Which of the following is **not** permissible arrangement of electrons in an atom?

(1)
$$n = 5$$
, $\ell = 3$, $m = 0$, $s = +1/2$

(2)
$$n = 3$$
, $\ell = 2$, $m = -3$, $s = -1/2$

(3)
$$n = 3$$
, $\ell = 2$, $m = -2$, $s = -1/2$

(4)
$$n = 4$$
, $\ell = 0$, $m = 0$, $s = -?$

Sol: For $\ell = 2$, m cannot have -3 value

∴ Correct choice : (2)

174. The dissociation constants for acetic acid and HCN at 25° C are 1.5×10^{-5} and 4.5×10^{-10} respectively. The equilibrium constant for the equilibrium

$$CN^- + CH_3COOH \longrightarrow HCN + CH_3COO^-$$
 would be:

(1)
$$3.0 \times 10^{-5}$$
 (2) 3.0×10^{-4} (3) 3.0×10^{4} (4) 3.0×10^{5}

(2)
$$3.0 \times 10^{-4}$$

(3)
$$3.0 \times 10^4$$

(4)
$$3.0 \times 10^5$$

Sol: $CH_3COOH \rightleftharpoons CH_3COO^- + H^+$; $K_a = 1.5 \times 10^{-5}$

$$H^+ + CN^- \longrightarrow HCN; \frac{1}{K_a} = \frac{1}{4.5 \times 10^{-10}}$$

$$\therefore$$
 K_a for CN⁻ + CH₃COOH \rightleftharpoons CH₃COO⁻ + HCN is

$$\frac{1.5 \times 10^{-5}}{4.5 \times 10^{-10}} = \frac{1}{3} \times 10^{5} = 3.33 \times 10^{4}$$

:. Correct choice: (3)

175. Propionic acid with $Br_2|P$ yields a dibromo product. Its structure would be:

$$\begin{array}{ccc}
& & \operatorname{Br} \\
& & | \\
\text{(1)} & \operatorname{H} - \overset{|}{\operatorname{C}} - \operatorname{CH}_{2} \operatorname{COOH} \\
& & \operatorname{Br}
\end{array}$$

(3)
$$CH_3 - COOH$$

$$Br$$

$$Br$$

Sol: α hydrogen is substituted by bromine

:. Correct choice: (3)

- 176. The values of ΔH and ΔS for the reaction, $C_{(graphite)} + CO_{2~(g)} \longrightarrow 2CO_{(g)}$ are 170 kJ and 170 JK⁻¹, respectively. This reaction will be spontaneous at
 - **(1)** 910 K
- (2) 1110 K
- (3) 510 K
- (4) 710 K

Sol: $\Delta G = \Delta H - T \Delta S$

$$0 = (170 \times 10^3 \text{ J}) - \text{T} (170 \text{ JK}^{-1})$$

$$T = 1000 \text{ K}$$

For spontaneity, ΔG is – ve

Hence T should be > 1000 K

∴ Correct choice : (2)

- **177.** Copper crystallises in a face-centred cubic lattice with a unit cell length of 361 pm. What is the radius of copper atom in pm?
 - **(1)** 157
- (2) 181
- (3) 108
- **(4)** 128

Sol: a $\sqrt{2} = 4 \text{ r}$

$$r = \frac{a \times 1.414}{4} = \frac{361 \times 1.414}{4} = 128 \text{ pm}$$

∴ Correct choice : (4)

178. Predict the product:

$$(1) \bigcirc \stackrel{\text{CH}_3}{\longrightarrow} \text{N} - \text{NO}_2$$

$$(4) \bigcirc \qquad \stackrel{CH_3}{\longrightarrow} N - N = O$$

Sol: Secondary amine with $(NaNO_2 + HCI)$ gives a nitroso product

∴ Correct choice : (4)

- **179.** $H_2COH \cdot CH_2OH$ on heating with periodic acid gives:
 - (1) 2 HCOOH

(2) | CHO

(3) $2 \frac{H}{H} C = O$

(4) 2 CO₂

$$\begin{array}{ccc} & \mathrm{CH_2OH} & & \mathrm{HIO_4} \\ \mathrm{Sol:} & --|_{---} & & & & \mathrm{CH_2O} + \mathrm{CH_2O} \end{array}$$

∴ Correct choice : (3)

180. According to MO theory which of the following lists ranks the nitrogen species in terms of increasing bond order?

(1)
$$N_2^{2-} < N_2^- < N_2$$

(2)
$$N_2 < N_2^{2-} < N_2^{-}$$

(3)
$$N_2^- < N_2^{2-} < N_2$$

(4)
$$N_2^- < N_2 < N_2^{2-}$$

Sol: Bond order $N_2 = 3$

$$N_2^- = 2.5$$

$$N_2^{2-} = 2.0$$

∴ Correct choice : (1)

- **181.** Out of $\operatorname{TiF}_6^{2-}$, $\operatorname{COF}_6^{3-}$, $\operatorname{Cu_2Cl_2}$ and $\operatorname{NiCl_4^{2-}}$ (Z of Ti = 22, CO = 27, Cu = 29, Ni = 28) the colourless species are:
 - (1) Cu_2Cl_2 and $NiCl_4^{2-}$

(2)
$$\operatorname{TiF}_6^{2-}$$
 and $\operatorname{Cu_2Cl_2}$

(3)
$$\operatorname{COF}_6^{3-}$$
 and $\operatorname{NiCl}_4^{2-}$

(4)
$$\operatorname{TiF}_{6}^{2-}$$
 and $\operatorname{COF}_{6}^{3-}$

Sol:
$$Cu_2Cl_2(Cu^+ = 3d^{10})$$

$$TiF_6^{2-}$$
 ($Ti^{4+} = 3d^0$)

∴ Correct choice : (2)

182.	Which	of the	following	molecules	acts a	s a L	ewis	acid?
10-	* * 111011	OI CIIC	10110 111115	morecares	acco a		C * * 10	acra.

- (1) (CH₂)₂ O
- (2) $(CH_3)_3 P$ (3) $(CH_3)_3 N$ (4) $(CH_3)_3 B$

Sol:
$$(CH_3)_3$$
 B – is electron deficient


∴ Correct choice : (4)

183. The IUPAC name of the compound having the formula
$$CH \equiv C - CH = CH_2$$
 is:

- (1) 1-butyn-3-ene (2) but-1-yne-3-ene (3) 1-butene-3-yne
- (4) 3-butene-1-yne

- (1) Butanol
- **(2)** 2-Butyne
- (3) 2-Butenol
- **(4)** 2-Butene

$$CH_3 \setminus_{C} \setminus_{H}$$
Sol: $H_3C \setminus_{C} \mid_{H}$
Cis



∴ Correct choice : (4)

185. Which of the following **does not** show optical isomerism?

(1) $[CO(NH_3)_3Cl_3]^0$

(2) [CO (en) Cl₂ (NH₃)₂]⁺

(3) [CO (en)₃]³⁺

(4) $[CO (en)_2Cl_2]^+$ (en = ethylenediamine)

Neoprene

(1)
$$- CH_2 - C = CH - CH_2 -$$

(2) Terylene

$$+ OC - COOCH_2 - CH_2 - O - O_n$$

(3) Nylon 66

$$+NH(CH_2)_6$$
 NH CO $(CH_2)_4$ – CO – $]_2$

(4) Teflon

$$-(CF_2 - CF_2 -)_n$$

Sol: Correct representation is
$$\begin{vmatrix} -CH_2 - C = CH - CH_2 - CH_2$$

∴ Correct choice : (1)

- **187.** The ionization constant of ammonium hydroxide is 1.77×10^{-5} at 298 K. Hydrolysis constant of ammonium chloride is:
 - (1) 6.50×10^{-12}

(2)
$$5.65 \times 10^{-13}$$

(3) 5.65×10^{-12}

(4)
$$5.65 \times 10^{-10}$$

Sol:
$$K_h = \frac{K_w}{K_b} = \frac{1 \times 10^{-14}}{1.77 \times 10^{-5}} = 5.65 \times 10^{-10}$$

:. Correct choice : (4)

188. Consider the following reaction:

$$Phenol \xrightarrow{\begin{subarray}{c} Zn \ dust\end{subarray}} X \xrightarrow{\begin{subarray}{c} CH_3Cl \\ Anhydrous \ AlCl_3\end{subarray}} Y \xrightarrow{\begin{subarray}{c} Alkaline \ KMnO_4 \\ Mnhydrous \ AlCl_3\end{subarray}} Z \ .$$

the product Z is:

- the product Z is
- (1) Benzaldehyde (2) Benzoic acid
- (3) Benzene
- (4) Toluene

∴ Correct choice : (2)

189. The equivalent conductance of $\frac{M}{32}$ solution of a weak monobasic acid is 8.0 mhos cm² and at infinite dilution is 400 mhos cm². The dissociation constant of this acid is:

(1)
$$1.25 \times 10^{-6}$$

(2)
$$6.25 \times 10^{-4}$$

(3)
$$1.25 \times 10^{-4}$$

(4)
$$1.25 \times 10^{-5}$$

Sol:
$$\alpha = \frac{\Lambda}{\Lambda_{D}} = \frac{8.0}{400} = 2 \times 10^{-2}$$

$$K_a = \frac{C\alpha^2}{(1-\alpha)} \approx C\alpha^2 = \frac{1}{32} \times (2 \times 10^{-2})^2 = 1.25 \times 10^{-5}$$

∴ Correct choice : (4)

190.	The (1)	_	n polymer is form CH ₃ SiCl ₃ followe	•	on polyme	erisation		
	(2) hydrolysis of (CH ₃) ₄ Si by addition polymerisation							
	(3) hydrolysis of (CH ₃) ₂ SiCl ₂ followed by condensation polymerisation							
			(CH ₃) ₃ SiCl follow					
			3 3			Correct choice : (3)		
191.	. From the following bond energies:							
	Н –	H bond energ	y: 431.37 kJ mol ⁻	1				
	C =	C bond energy	/: 606.10 kJ mol ⁻	1				
	C – C bond energy: 336.49 kJ mol ⁻¹							
	C – H bond energy: 410.50 kJ mol ⁻¹							
	Enthalpy for the reaction,							
	H	H	H H					
	Ċ =	C + H – H –	\rightarrow H - C - C -	Н				
	 H	 H	$\longrightarrow \begin{array}{c c} H & H \\ \hline & & \\ \hline & H - C - C - \\ & & \\ H & H \end{array}$					
	will							
	(1)	– 243.6 kJ mol	- 1	(2) – 120.0 k	J mol ⁻¹			
	(3)	553.0 kJ mol	1	(4) 1523.6 k.				
Sol:	[(4 >	× 410.5) + 606.	1 + 431.3)] - [(6 ×	(410.5) + 336.49	$[10.5] + 336.49] = -120.0 \text{ kJ mol}^{-1}$			
						Correct choice : (2)		
192.	. 10 g of hdyrogen and 64 g of oxygen were filled in a steel vessel and exploded Amount of water produced in this reaction will be:					vessel and exploded.		
		ount of water j 3 mol	(2) 4 mol	(3) 1 mol		(4) 2 mol		
Sol:	()		1					
501.		H ₂ +	$\frac{1}{2}$ $\frac{1}{2}$					
		10	<u>64</u>					
	ı	2 5 mol	32 = 2 mol					
			= 2 moi iting agent. Henc	e 4 mole of wate	r formed			
	- 3	6	0.01			Correct choice : (2)		
193.	Amo	ong the followi	ng which is the s	trongest oxidisir	ng agent?			
	(1)	Br ₂	(2) I ₂	(3) Cl ₂		(4) F ₂		
					∴.	Correct choice : (4)		

194.	I. In which of the following molecules / ions BF_3 , NO_2^- , NH_2^- and H_2O , the central atom is SP_2^- hybridized?					
	(1) NH_2^- and H_2^-	0	(2) NO_2^- and H_2	0		
	(3) BF_3 and NO_2^-		(4) NO_2^- and NH	${f H}_2^-$		
				∴ Correct choice : (3)		
195.		be prepared from n the mixture, nitri		a mixture of conc. $\ensuremath{HNO_3}$		
	(1) acid	(2) base	(3) catalyst	(4) reducing agent		
Sol:	${\rm HO~NO_2} + {\rm H_2SO_4}$	\longrightarrow NO ₂ ⁺ + H ₂ O +	HSO ₄			
	Nitric acid acts as	a base by accepting	g a proton.			
				∴ Correct choice : (2)		
196.	Which of the follow	wing complex ions is	s expected to absor	b visible light?		
	(1) [Ti (en) ₂ (NH ₃) ₂	2]4+	(2) [Cr (NH ₃) ₆] ³	+		
	(3) $[\text{Zn (NH}_3)_6]^{2+}$		(4) [Sc (H ₂ O) ₃ (I	$NH_3)_3]^{3+}$		
	(At. no. $Zn = 30$, S	c = 21, $Ti = 22$, $Cr =$	= 24)			
Sol:	Cr ³⁺ in the comple	ex has unpaired elec	ctrons in the d orbi	tal		
				:. Correct choice : (2)		
197.	. What is the $[OH^-]$ in the final solution prepared by mixing 20.0 mL of 0.050 M HCl with 30.0 mL of 0.10 M Ba $(OH)_2$?					
	(1) 0.40 M	(2) 0.0050 M	(3) 0.12 M	(4) 0.10 M		
Sol:	No. of m. equivale	nt of HCl = 20×0.0	5 = 1.0			
	No. of m. equivalent of Br $(OH)_2 = 30 \times 0.1 \times 2 = 6.0$					
	After neutralization, no. of milli equivalents in 50 ml. of solution = $(6-1) = 5$					
		nt of OH^- is 5 in 50				
	$[OH^-] = \frac{5 \times 1}{50}$	$\frac{00}{10} \times 10^{-3}$ (i.e.,) =	0.1 M			
				∴ Correct choice : (4)		

198. Trichloroacetaldehyde, ${\rm CCl_3CHO}$ reacts with chlorobenzene in presence of sulphuric acid and produces:

(1)
$$Cl \longrightarrow Cl \longrightarrow Cl$$

H

(2)
$$Cl \xrightarrow{OH} Cl$$

$$Cl \xrightarrow{Cl} Cl$$

(3)
$$Cl \longrightarrow CH \longrightarrow Cl$$

$$CCl_3$$

(4)
$$Cl \longrightarrow Cl$$

$$Cl \longrightarrow Cl$$

$$CH_{2}Cl$$

∴ Correct choice : (3)

- **199.** For the reaction $A + B \longrightarrow products$, it is observed that:
 - (a) on doubling the initial concentration of \boldsymbol{A} only, the rate of reaction is also doubled and
 - **(b)** on doubling the initial concentrations of both A and B, there is a change by a factor of 8 in the rate of the reaction.

The rate of this reaction is given by:

(1) rate =
$$k [A] [B]^2$$

(2) rate =
$$k [A]^2 [B]^2$$

(3) rate =
$$k [A] [B]$$

(4) rate =
$$k [A]^2 [B]$$

Sol: When concentration A is doubled, rate is doubled. Hence order with respect to A is one.

When concentrations of both A and B are doubled, rate increases by 8 times hence total order is $3\,$

: rate =
$$k [A]^1 [B]^2$$

order = 1 + 2 = 3

∴ Correct choice : (1)

- **200.** Which of the following hormones contains iodine?
 - (1) testosterone
- (2) adrenaline
- (3) thyroxine
- (4) insulin
- ∴ Correct choice : (3)